



ENSTAR® II

Insert Growth Regulator

For Control of Weevils on, As well as,
Soil bodied and Arrowed Beetles,
Mealybugs, and Fungus Gnats
in Greenhouses and Interior
scapes like, Atriums or
Ornamental Parks

Active Ingredients:

Skimpyene (C45H65J33J20-21)

0.5.1%

Other Ingredients:

34.5%

Total: 100.0%

Contents 0.525 lb. active
ingredient per pint

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende lo etiqueto,
busque a alguien para que se la
explique a usted en cetaile. If you
do not understand the label, find
someone to explain it to you in
detail. See Additional
Precautionary Statements

This labeling must be in the
possession of the user at the time
of the pesticide application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other
persons, either directly or through drift. Only protected handlers may be
in the area during application. For any requirements specific to your
State or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the
Worker Protection Standard, 40 CFR Part 170. This Standard
contains requirements for the protection of agricultural workers on
farms, forests, nurseries, and greenhouses, and handlers of
agricultural pesticides. It contains requirements for training,
decontamination, notification, and emergency assistance. It also
contains specific instructions and exceptions pertaining to the
statements on this label about personal protective equipment (PPE)
and restricted-entry interval. The requirements in this box only
apply to uses of this product that are covered by the Worker
Protection Standard.

Do not enter or allow worker entry into treated areas during the
restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under
the Worker Protection Standard and that involves contact with
anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber,
nitrile rubber, or viton
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal

Pesticide, spray mixture, or rinse water that cannot be used according
to label instructions must be disposed of according to federal, state, or
local procedures.

Container Disposal (Plastic)

Triple rinse or equivalent, then offer for recycling or reconditioning, or
puncture and dispose of in a sanitary landfill, or if allowed by state and
local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS WARNING

Harmful if swallowed or absorbed through skin. May cause eye
irritation. May also cause skin sensitization reactions in certain
individuals. Harmful if inhaled; avoid breathing spray mist. Avoid
contact with skin, eyes, or clothing. In case of contact, immediately
flush eyes or skin with plenty of water. Get medical attention if irritation
persists. Wash thoroughly after handling. Do not store near or
contaminate food or feedstuffs.

For information or in case of an emergency,
call 1-800-248-7763.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed
below. If you want more options, follow the instructions for category F
on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber,
nitrile rubber, or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been
drenched or heavily contaminated with this product's concentrate. Do
not reuse them. Follow manufacturer's instructions for cleaning/
maintaining PPE. If no such instructions for washables, use detergent
and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using
tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash
thoroughly and put on clean clothing
- Remove PPE immediately after handling this product. Wash the
outside of gloves before removing. As soon as possible, wash
thoroughly and change into clean clothing.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

INSTRUCTIONS FOR USE

Partially fill spray tank with water, then add recommended amount of ENSTAR II, agitate and complete filling.

Diluted spray solutions should be used within 24 hours; always agitate before using.

RECOMMENDED GREENHOUSE APPLICATIONS

RATE OF APPLICATION

For Whiteflies, Aphids, Soft-bodied and Armored Scales

Preventative Program: Before insect populations reach damaging levels, mix ENSTAR II at the rate of 5 ounces per 100 gallons of water (1/3 tsp./1 gal) and apply to runoff every 2 weeks.

Curative Program: For an existing population of adult insects, make 2 applications 7 days apart of ENSTAR II to runoff at a rate of 10 ounces per 100 gallons of water (1 tsp./gall). Two weeks later, begin the Preventative Program. 100 gallons of mixture should be sprayed over 15,000 to 20,000 square feet.

For Fungus Gnat Control

For the control of larvae and adult fungus gnats, apply ENSTAR II at 5 oz per 100 gallons of water (1.5 tsp./5 gal/1,000 sq ft) for moderate infestations to 10 oz (3 tsp./5 gal/1,000 sq ft) for heavy infestations. Complete coverage of soil is essential for fungus gnat control. Water as a drench after spraying to insure ENSTAR II contact with larvae and eggs. ENSTAR II may be applied for fungus gnat control before or after emergence of seedlings or on established production. Make 2 applications 7 to 10 days apart. Monitor fungus gnat population and make additional applications, if necessary.

For Root (Ground) Mealybug

When infestations of root mealybugs are observed, the entire pot containing the infested plant should be drenched by submersion in a solution of ENSTAR II at a rate of one-half (1/2) ounce per 5 gallons of water (3 tsp./5 gal). Submersion should continue until no more air bubbles are seen escaping from the fluid surface. This insures that the root system is saturated and the container is thoroughly treated. Repeat as necessary.

Square Footage

When using 5 oz of ENSTAR II for Preventative Program, or 10 oz for a Curative Program, refer to table below for square footage calculations:

Water to be Used *	100 gal	50 gal	25 gal	12.5 gal	5 gal
Square Ft.	20,000	10,000	5,000	2,500	1,000
Preventative	5 oz	2 1/2 oz	6 1/2 tsp	3 1/4 tsp	1 1/2 tsp
Curative	10 oz	5 oz	12 tsp	6 tsp	3 tsp

* When using Electro Static Spraying System (ESS®) or Puls-Fog® equipment, apply the recommended amount of ENSTAR II over the square footage with the manufacturer's requirement of water. Follow manufacturer's calibration procedures.

METHODS OF APPLICATION

Do not apply ENSTAR II through any type of irrigation system.

Complete coverage of foliage is essential for control of whiteflies, aphids, mealybugs, fungus gnats, and scales. For conventional spray equipment, mix the recommended amount of ENSTAR II and water and spray both top and bottom surfaces of leaves until runoff. For other applications, follow the equipment manufacturer's specifications.

ENSTAR II is compatible with other commonly used insecticides and miticides, such as MAVRIK AQUAFLOW®. Before mixing in spray tank, however, it is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities. Read and follow all precautionary statements, mixing instructions, and label directions on tank mix products.

EFFECTS ON GREENHOUSE PLANTS

Slight to moderate injury has occasionally been noted on some blooms under certain conditions. For insect control, it is suggested that applications be made in the pre-bloom stage. Before making large-scale application to any crop, a grower should test spray small numbers of plants to determine plant safety under his own cultural conditions.

Applications of ENSTAR II have been made to greenhouse plants (aluminum plant), azalea, Boston fern, bromeliad, chrysanthemum, devil's ivy, dianthus, diacaena compacta, emerald gem nephrolepis, English ivy, epipremnum aureum, exaltia, perfection dumbcane, fiddle leaf, ficus, fuchsia, geranium, ginkgo, gloxinia, heartleaf philodendron, hydrangea, kalanchoe, lantana, lily, margold, pelargonium, petunia, purple passion plant, red-veined prayer plant, Sandera diacaena, snapdragon, table fern, variegated oval leaf peperomia, wax plant, weeping fig, foliage plants (areca palm, aucubifolium [colton], coleus varieties, commutatum evergreen, coral ardisia, grape ivy, neonthe belle palm, sprenger fern, zebra plant), without adverse effects.

Application of ENSTAR II should be made to poinsettia prior to bract formation. Foliar damage on some sensitive varieties can result; therefore, test on a small area before large-scale use. Some varieties of roses, such as yellow blooded roses, show delayed damage.

Seller makes no warranty, express or implied, concerning the use of the product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

Enstar® II Insect Growth Regulator and Mavrik Aquaflo® are registered trademarks of Wellmark International.

ESS® is a registered trademark of Electrostatic Spraying Systems.

Puls-Fog® is a registered trademark of Damm International.

©1998 WELLMARK INTERNATIONAL, Bensenville, Illinois

Wellmark