

# Specimen Label



# ZeroTol<sup>TM</sup>

Broad Spectrum Algaecide/Fungicide

**PROFESSIONAL TURF CARE**



**KEEP OUT OF REACH OF CHILDREN**  
**ANGER - PELIGRO**

**ACTIVE INGREDIENT:**

Hydrogen Dioxide.....27%  
INERT INGREDIENTS .....73%  
TOTAL.....100%

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

EPA Registration No. 70299-1

EPA Establishment No. 60156-IL-001

Sold by: **Bi Safe Systems**  
Glastonbury, CT 06033

- \* Preventative treatment for ornamental plants and turf.
- \* Professional Turf Care.

## FOR HORTICULTURAL AND COMMERCIAL USE ONLY

### FIRST AID

#### If in eyes

- Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 – 20 minutes.
- Call a poison control center or doctor for treatment advice.

#### If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center.
- Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

#### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

**Sold by:** BioSafe Systems, 80 Commerce Street,  
Glastonbury, CT 06033

**EPA Registration No.** 70299-1

**EPA Establishment No.** 60156-IL-001

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

**CORROSIVE:** Concentrate causes irreversible eye damage. Concentrate may be fatal if swallowed. Concentrate causes skin irritation or temporary discoloration on exposed skin. Do not breathe vapor of concentrate. Do not get concentrate in eyes, on skin or on clothing.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

When handling concentrate wear protective eyewear (goggles or face shield) and rubber gloves. Applicators and handlers must wear coveralls over long-sleeved shirt, long pants, and chemical resistant footwear plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water.

#### USER SAFETY RECOMMENDATIONS

Users should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

**FOR TERRESTRIAL USES.** Keep out of lakes, ponds and streams. This pesticide is toxic to birds and fish. Do not apply directly to water, or to areas where surface water is present or to inter-tidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wash waters.

This product is highly toxic to bees and other beneficial insects exposed to direct contact on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Do not apply this product or allow it to drift to crops where beneficials are part of an Integrated Pest Management strategy.

### PHYSICAL AND CHEMICAL HAZARDS

Strong oxidizing agent. **Corrosive.** Do not use in concentrated form. Mix only with water in accordance with label instructions. Never bring concentrate in contact with other pesticides, cleaners or oxidative agents.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted-Entry Interval (REI). The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

**THERE IS A RESTRICTED ENTRY OF ZERO (0) HOURS FOR THIS PRODUCT.**

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Store in original containers in a cool, well-ventilated area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited. If waste cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or dispose in a sanitary landfill, or incineration, if allowed by state and local authorities by burning, stay out of smoke.

ZeroTol works by surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted. ZeroTol does not produce any visible residue, distinct odor or deleterious effects to plants when used in accordance with label directions.

#### Compatibility:

Do not use at higher than recommended dilution rates as leaf burn may result. ZeroTol has been designed to provide a balanced source of the active ingredient directly to the plant surface and has been shown to not cause adverse cosmetic effects on most plants. Since we have not tested all plant species, however, it is always advisable to test ZeroTol on a few plants before treating large numbers.

#### Solution Preparation:

\* ZeroTol works best when diluted with water containing low levels of organic or inorganic materials and having a neutral pH. Thoroughly rinse out mixing tank with water before mixing concentrate. ZeroTol will readily mix with clean, neutral water and does not require agitation.

\* ZeroTol concentrate should not be combined or mixed with any other pesticide or fertilizer.

\* ZeroTol is formulated with minimal surfactant for plants having waxy or hairy surfaces. Additional surfactant may be added, if needed for treatment of plants with difficult to reach surfaces.

\* ZeroTol is a strong oxidizing agent and may react with residues of metal-based fungicides or supplements. Care should be used when applying ZeroTol as a foliar spray immediately following foliar applications of metal-based products.

#### For turf applications:

Broad spectrum treatment for control of algae, fungi and bacteria on turf.

For use on all turf types such as commercial turf, lawns, athletic fields and golf course fairways, greens and tees.

Use ZeroTol to control fungi such as: *Anthracnose*, Brown Spot, Dollar Spot, Copper Spot, Fairy Ring, Pink Snow Mold, *Pythium*, Phytophthora, Summer Patch, Rhizoctonia, Scum, Take All Patch *Fusarium* Blight, Stripe Smut, Leaf Spot, Algae, Slime Molds and their spores.

ZEROTOL CONTROLS ON CONTACT.

**For treatment of turf,** use on golf course fairways, greens and tees of Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass and their mixtures to control/suppress algae, bacterial and fungal diseases, and the odors and conditions that these organisms may cause.

Typical treatment rates involve treating approximately 1000 square feet of lawn area with 3 to 10 gallons of diluted solution. Add a spreader surfactant for best results. **Refer to Table 1. for turf application rates.**

- Optimum treatment time is early morning or late afternoon.
- For best results, apply immediately after grass has been cut.
- Applications can be made during wet or rainy weather.
- Use spray solution the same day it is prepared, do not store and reuse mixed spray solution.
- ZeroTol can be injected through automatic irrigation systems in turf areas. Refer to Chemigation Directions for Use for specific instructions on using this product through irrigation systems.

**For seed bed treatment** – Prior to sowing seed, use a dilution of 1:50 or 2½ fl. oz. per gallon of clean water. Thoroughly wet or drench the seedbed, to the point of saturation, with 60 to 100 gallons of dilute solution per 1000 square feet. Let sit for one hour then immediately seed soil.

After seeds have germinated, use a dilution of 1:100 or 1¼ fl. oz. per gallon of clean water. Lightly spray or irrigate the soil and seedlings until thoroughly wetted. Retreat once per week until seed is well established.

**For soil treatment, pre-inoculation with beneficial organisms,** use ZeroTol to reduce the number of potentially plant pathogenic organisms in the soil that will prevent beneficials from becoming established. Use a dilution of 1:50 or 2½ fl. oz. per gallon of clean water. Thoroughly wet or drench the area to be inoculated. Wait one day before inoculating soil.

**For turf treatment, post-inoculation with beneficial organisms,** use ZeroTol to control plant pathogens on the foliar portion of turf. Careful application of ZeroTol must be made to prevent drenching into the root system and subsequent temporary reduction in soil organisms.

Use a dilution of 1:100 or 1¼ fl. oz. per gallon of clean water. Apply to the turf by lightly spraying leaf surfaces. Do not allow

solution to be drenched into the soil and root system. Drenching of ZeroTol into the soil will result in temporary reduction of beneficial organism colonies that are directly affected.

**CHEMIGATION DIRECTIONS FOR USE**  
**General Requirements:**

- 1) Apply this product only through a sprinkler including a center pivot, lateral move, end tow, side wheel roll, traveler, solid set, hand move, flood basin, humidification or drip trickle irrigation system, or through misting systems.
- 2) Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- 3) Ensure that the irrigation system used is properly calibrated and if you have questions, call the state extension service or the equipment manufacturer.
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless proper safety devices for public water systems are in place. Read label for instructions.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.

**Specific Requirements:**

- 1) Public water supply means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of 25 individuals daily at least 60 days throughout the year.
- 2) Chemigation systems connected to the public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of the overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of liquid back towards the injector.
- 4) The pesticide injection pipeline must contain a functional, normally closed, solenoid, operated valve located on the

intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.

- 5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump, or equivalent, effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment.

**Application Instructions:**

- 1) Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues may cause product to loose effectiveness or strength.
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions.
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required. The product will immediately go into suspension without any required agitation.
- 4) ZeroTol should not be applied in conjunction with any other pesticides or fertilizers; this may cause reduced performance of the product and should be avoided.

**WARRANTY**

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing, method of application, weather, watering practices, nature of soil, potting medium, disease problem, condition of crop, incompatibility with other chemicals, pre-existing conditions and other conditions influencing the use of this product are beyond the control of the seller. Buyer assumes all risks associated with the use, storage, or handling of this material not in strict accordance with directions given herewith. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.

**Table 1.**

Disease Controlled	Curative Rate	Preventative Rate	Notes
<b>Anthracnose</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle.Combine with a systemic fungicide for residual suppression.
<b>Brown Spot</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle.Combine with a systemic fungicide for residual suppression.
<b>Dollar Spot</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.

Disease Controlled	Curative Rate	Preventative Rate	Notes
<b>Copper Spot</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.
<b>Summer Patch</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.
<b>Stripe Smut</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.
<b>Take All Patch</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.
<b>Leaf Spot</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.
<b><i>Fusarium Blight</i></b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Once control is achieved, follow with a 7-day prevention cycle. Combine with a systemic fungicide for residual suppression.
<b>Fairy Ring</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Drench the soil to saturate the root systems in areas affected. Use 5 - 10 gallons per 1000 sq. ft.
<b>Pink Snow Mold</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Spray in early fall to reduce number of dormant spores. Treat throughout winter. May be applied to frozen ground.
<b><i>Pythium</i></b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Drench the soil to saturate the root systems in areas affected. Use 5 - 10 gallons per 1000 sq. ft.
<b><i>Phytophthora</i></b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Drench the soil to saturate the root systems in areas affected. Use 5 - 10 gallons per 1000 sq. ft.
<b><i>Rhizoctonia</i></b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Drench the soil to saturate the root systems in areas affected. Use 5 - 10 gallons per 1000 sq. ft.
<b>Algae &amp; Slime Molds, Scum</b>	6 – 12 fl. oz. per 1000 sq. ft. Use 3 – 5 gallons of solution per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Drench the soil to saturate the root systems in areas affected. Use 5 - 10 gallons per 1000 sq. ft.
<b>Heavy Algae</b>	12 – 25 fl. oz. per 1000 sq. ft.	2 – 6 fl. oz. per 1000 sq. ft. Apply at 7-day intervals.	Curative control may require 2 to 3 consecutive treatments to eradicate disease. Drench the soil to saturate the root systems in areas affected. Use 5 - 10 gallons per 1000 sq. ft.